

Global Observables at PHENIX

Alexander Bazilevsky¹, for the PHENIX Collaboration

¹ RIKEN BNL Research Center, Brookhaven National Laboratory,
Upton, NY, USA

Abstract. We present PHENIX recent results on charged particle and transverse energy densities measured at mid-rapidity in Au-Au collisions at $\sqrt{s_{NN}}=130$ GeV and 200 GeV over broad range of centralities. The mean transverse energy per charged particle is derived. The first PHENIX measurements at $\sqrt{s_{NN}}=19.6$ GeV are also presented. A comparison with calculations from various theoretical models is performed.

Keywords: heavy ion, nuclear dynamics, particle multiplicity, transverse energy
PACS: 25.75.-q

1. Introduction

Charged particle multiplicity (N_{ch}) and transverse energy (E_T) are global variables which give excellent characterization of the high energy nucleus-nucleus collisions, thus providing information about the initial conditions [1]. They help to constrain the wide range of theoretical predictions and discriminate among various mechanisms of particle production. N_{ch} measurements also provide one with an opportunity to study high density QCD effects in relativistic nuclear collisions [2].

First results for N_{ch} and E_T at mid-rapidity in Au-Au collisions at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV measured with the PHENIX detector were published in [3, 4, 5, 6]. In this paper we extend the N_{ch} and E_T analysis to more peripheral collisions, up to 65%-70% centrality class corresponding to the average number of participating nucleons $N_p \approx 22$. The results for the $\langle E_T \rangle$ / $\langle N_{ch} \rangle$ are presented at $\sqrt{s_{NN}} = 19.6$ GeV, 130 GeV and 200 GeV.

We used the same experimental techniques to analyze data obtained at different beam energies. The minimum bias (MB) trigger in $\sqrt{s_{NN}} = 19.6$ GeV data was formed by the coincidence of two beam-beam counters. In $\sqrt{s_{NN}} = 130$ GeV and 200 GeV data the MB trigger additionally required the signals from two zero degree calorimeters [3, 4, 5, 6]. The corrections for the measured E_T and N_{ch} related to particle composition and mean transverse momentum are performed based on PHENIX results obtained at $\sqrt{s_{NN}} = 130$ GeV [7, 8] and 200 GeV [9]. $\sqrt{s_{NN}} =$




Fig. 1. $dE_T/d\eta$ (top panels) and $dN_{ch}/d\eta$ (bottom panels) per pair of participants versus N_p (top) and $dN_{ch}/d\eta$ (bottom) versus centrality; N_p is taken from data at $\sqrt{s_{NN}} = 130$ GeV (left panels) and $\sqrt{s_{NN}} = 200$ GeV (right panels); The lines represent the effect of the $\pm 1\sigma$ centrality-dependent systematic errors, the error bars are the total systematic errors.

Fig. 2. $R_{200/130}$ for $dE_T/d\eta$ (top) and $dN_{ch}/d\eta$ (bottom) versus centrality; N_p is taken from data at $\sqrt{s_{NN}} = 200$ GeV; for the explanation of error representation, see the caption of Fig. 1.

19.6 GeV data corrections are based on NA49 measurements obtained at $\sqrt{s_{NN}} = 17.2$ GeV [10].

The part of the PbSc electromagnetic calorimeter used for the E_T measurements covers the pseudorapidity range $|\eta| \leq 0.38$ with an azimuthal aperture of $\Delta\phi = 44.4^\circ$ in Run-1 data ($\sqrt{s_{NN}} = 130$ GeV) and $\Delta\phi = 112^\circ$ in Run-2 data ($\sqrt{s_{NN}} = 19.6$ GeV and 200 GeV). The pad chambers used for N_{ch} measurements have a fiducial aperture of $|\eta| \leq 0.35$ and $\Delta\phi = 90^\circ$ and 180° in the Run-1 and Run-2 data respectively.

2. Results

Fig. 1 shows the centrality dependence of $dE_T/d\eta$ and $dN_{ch}/d\eta$ per participant pair measured at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV. Both values show a steady rise with N_p .

The ratio of the pseudorapidity densities measured at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV ($R_{200/130}$) for each centrality bin, corresponding to 5% of the nuclear




Fig. 3. The comparison of $dN_{ch}/d\eta$ at $\sqrt{s_{NN}} = 130$ GeV (top panels) and 200 GeV (middle panels) along with $R_{200/130}$ (bottom panels) as a function of centrality to the following models (solid lines): HIJING [12], EKRT [11], KLN [2], mini-jet [14] (shaded area), AMPT [15] and String Fusion model [13].

interaction cross section, is shown in Fig. 2. The centrality dependence of the ratios is consistent with a constant from the most central to semiperipheral collisions. For $N_p < 70$, ratios tend to drop.

Fig. 3 shows the comparison of our results to different model predictions. A powerful test for theoretical models is the comparison to the measured ratios $R_{200/130}$, since many systematic errors in experimental measurements cancel out. The increase of $dN_{ch}/d\eta$ with centrality is in contrast to the predictions of the EKRT model [11]. HIJING [12] is in qualitative agreement with this scenario, however the strong centrality dependence of the ratio $R_{200/130}$ predicted by HIJING is excluded by the data. Demonstrating a very good agreement with PHENIX measurements at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV, String Fusion model [13] seems to underestimate the $R_{200/130}$. Our experimental results are well described by two-component mini-jet [14] and high energy QCD gluon saturation [2] model calculations from central to semiperipheral events. The PHENIX measurements of the ratio $R_{200/130}$ dropping at $N_p < 70$ might set a centrality limit for gluon saturation model application, which predicts a near constant behavior of $R_{200/130}$ vs N_p .




Fig. 4. $dE_T/d\eta|_{\eta=0} / dN_{ch}/d\eta|_{\eta=0}$ versus N_p

E_T and N_{ch} behave in a very similar manner such that the mean E_T per charged particle remains unchanged over a broad range of centralities (see Fig. 4). Fig. 4 also presents our preliminary result obtained at $\sqrt{s_{NN}} = 19.6$ GeV. Since we haven't yet finalized the centrality determination in $\sqrt{s_{NN}} = 19.6$ GeV data, we present our result with one point with large uncertainty in N_p (this error cancels out in the ratio $\langle E_T \rangle / \langle N_{ch} \rangle$).

3. Conclusions

We presented our results for N_{ch} and E_T in a broad range of centralities corresponding to N_p varying from 22 to 350. Our results were compared to the various theoretical model calculations. Two-component mini-jet [14] and high energy QCD gluon saturation [2] model calculations describe our data reasonably well. The ratio $R_{200/130}$ of charged particle multiplicities measured at $\sqrt{s_{NN}} = 200$ GeV and 130 GeV tends to drop at $N_p < 70$. This implies a centrality limit for gluon saturation model application, which predicts a near constant behavior of $R_{200/130}$.

The ratio of the mean E_T and N_{ch} stays near constant over the broad range of centralities. It also doesn't change from $\sqrt{s_{NN}} = 130$ GeV to 200 GeV within our systematic errors. However, the $\langle E_T \rangle / \langle N_{ch} \rangle$ measured at $\sqrt{s_{NN}} = 19.6$ GeV was about 20% below that obtained at higher $\sqrt{s_{NN}}$. This is consistent with mean p_t decrease from RHIC $\sqrt{s_{NN}} = 130$ GeV and 200 GeV [7, 9] to SPS $\sqrt{s_{NN}} = 17.2$ GeV [10].

Acknowledgements

We thank the staff of the RHIC project, Collider-Accelerator, and Physics Departments at BNL and the staff of PHENIX participating institutions for their vital contributions. We acknowledge support from the Department of Energy and NSF (U.S.A.), MEXT and JSPS (Japan), RAS, RMAE, and RMS (Russia), BMBF, DAAD, and AvH (Germany), VR and the Wallenberg Foundation (Sweden), MIST and NSERC (Canada), CNPq and FAPESP (Brazil), IN2P3/CNRS (France), DAE and DST (India), LG-YF, KRF and KOSEF (Korea), and the US-Israel Binational Science Foundation.

References

1. S.A. Bass et al., Nucl. Phys. **A661**, 205 (1999).
2. D. Kharzeev and M. Nardi, Phys. Lett. **B507**, 121 (2001); D. Kharzeev and E. Levin, Phys. Lett. **B523**, 79 (2001).
3. K. Adcox et al., Phys. Rev. Lett. **86**, 3500 (2001).
4. K. Adcox et al., Phys. Rev. Lett. **87**, 052301 (2001).
5. A. Milov, Nucl. Phys. **A698**, 171 (2002).
6. A. Bazilevsky, Nucl. Phys. **A715**, 486 (2003).
7. K. Adcox et al., Phys. Rev. Lett. **88**, 242301 (2002).
8. K. Adcox et al., Phys. Rev. Lett. **89**, 092302 (2002).
9. T. Chujo, nucl-ex/0209027.
10. J. Bächler et al., Nucl. Phys. **A661**, 45 (1999); T. Alber et al., Phys. Rev. Lett. **75**, 3814 (1995).
11. K.J. Eskola et al., Nucl. Phys. **B570**, 379 (2000); Phys. Lett. **B497**, 39 (2001).
12. X.N. Wang and M. Gyulassy, Phys. Rev. Lett. **86**, 3498 (2001).
13. N. Armesto et al., Phys. Lett. **B527**, 92 (2001); N.S. Amelin et al., Eur. Phys. J **C22**, 149 (2001).
14. S. Lee and X.N. Wang, Phys. Lett. **B527**, 85 (2002).
15. B. Zhang et al., Phys. Rev. **C61**, 067901 (2000).